

Effectiveness of the use of semiautomatic defibrillation in basic life support services

JOSÉ LUIS MORENO MARTÍN, ÓSCAR ESQUILAS SÁNCHEZ, ERVIGIO CORRAL TORRES, ROSA MARÍA SUÁREZ BUSTAMANTE, MARÍA ISABEL VARGAS ROMÁN

Servicio Municipal de Urgencia y Rescate del Ayuntamiento de Madrid (SAMUR-Protección Civil). Madrid, España.

CORRESPONDENCE:

Ervigio Corral Torres
Subdirección General SAMUR-
Protección Civil
Ronda de las provincias, s/n
28011 Madrid, España
E-mail: corralte@municipalidadmadrid.es

DATE OF RECEIPT:

4-9-2007

DATE OF ACCEPTANCE:

4-9-2008

CONFLICT OF INTEREST:

None

Objective: The objective of this study was to determine the importance of the use of semiautomatic defibrillation for the treatment of out-of-hospital cardiorespiratory arrest of any etiology attended initially by basic life support ambulant services. The analysis was based on survival and neurological evaluation 7 days after the event.

Design: A prospective, descriptive, transverse, longitudinal study that analyzed survival and the neurological status using the Glasgow-Pittsburgh scale 7 days after cardiorespiratory arrest in 233 patients attended over a 5-year period by a two-tier, urban, out-of-hospital emergency service based in Madrid, Spain (SAMUR-PC).

Results: Survival with complete recovery observed after out-of-hospital cardiorespiratory arrest was 7.29%, a figure very similar to the results for recovery in our department (12%), which included the arrests attended directly by advanced life support services. There was a significant difference between the patients given early defibrillation by the basic life support team and those in whom this was not performed based on the initial rhythm. It was also found that the initial call for assistance specified cardiorespiratory arrest or unconsciousness in only 37.7% of cases.

Conclusions: The primary response to cardiorespiratory arrest by basic life support ambulance services with semiautomatic external defibrillators can achieve survivals similar to those in patients attended directly by advanced life support services. [Emergencias 2009;21:12-16]

Key words: Basic life support. Semiautomatic defibrillation. Emergency techniques. Cardiorespiratory arrest. Survival.

Introduction

The management of cardiopulmonary arrest (CPA) occurring in public may constitute one of the main challenges faced by an Out-of-hospital Medical Emergency Service. Rapid processing of the call, correct assignment of resources, a short response time, adequate staff training in resuscitation techniques and evidenced-based procedures are some of the keys to the successful management of this pathology¹⁻¹¹. The relation between survival of these patients and response time is widely established^{12,13}. The search for new, useful and effective instruments and intervention methods should be permanent, since they may prove vital when faced with a case of CPA¹⁴. For public access defibrillation¹⁶, it is necessary to ensure correct area distribution of out-of-hospital attendance units in order to achieve early arrival at the scene of the event⁹. This is clearly not possible using only medicalised resources [Advanced Life

Support (ALS) Units]. In a two-tier Out-of-hospital Emergency Service such as SAMUR Protección Civil (SAMUR-PC), whose most numerous resources are Basic Life Support (BLS) units staffed by two emergency technicians and equipped for defibrillation, with a second tier of ALS units, staffed by a physician, qualified nurse and emergency technician, it is necessary to discriminate exactly what type of resource should be sent out. To suppose that the first response to a CPA should "always and only" be an ALS unit is today a clearly inefficient measure in health-care management¹⁷⁻¹⁹. Professional training of emergency technicians, together with adequately equipping BLS units, are important to shorten the time taken to provide initial care of the^{3-6,20-24}.

In 1995, SAMUR-PC incorporated the first semiautomatic defibrillators (AED) into the service²⁵ and today all BLS units carry this device. The main objective of this study was to evaluate the effectiveness of AED implementation in the BLS units of

an Out-of-hospital Medical Emergency Service. We wished to determine survival data of those patients initially defibrillated by these units versus those who were not, or those initially defibrillated by ALS units.

Method

A prospective, descriptive, longitudinal, cross-sectional study was designed. The study setting was the out-of-hospital medical emergency service of the Madrid City Council (SAMUR-PC). This service is organized as a two-tier system, with BLS units equipped with AED and ALS units. It is activated by calls to the single European emergency number (112) and the criteria of response to demands were established as a function of severity and the time taken to reach the patient. SAMUR-PC has a rigorous quality control system that includes, among other measures, follow-up of all critical patients at 6 hours, 24 hours and 7 days after admission to hospital.

The study population included all those patients who suffered an out-of-hospital CPA and were first attended by a BLS unit equipped with a AED during a 5-year period from February 2001 to February 2006. Variables studied were: age, sex, cause of initial activation, response time (from the end of the call requiring attendance to the moment of attendance), survival at 6 hours, 24 hours and at 7 days, and cerebral function evaluation using the Glasgow-Pittsburg Scale at 7 days after admission. Statistical analysis of data stored on Access 2000 was performed using SPSS v.13.0. Qualitative variables are described as frequencies and quantitative variables as central means and dispersion. Qualitative variables were compared using chi square test and means were compared using Student t test and non-parametric tests (Mann-Whitney U) for non-normal distributions. A p value of < 0.05 was considered statistically significant. Ethical aspects were respected and data confidentiality guaranteed.

Results

The criteria for inclusion were met by 233 patients (Figure 1). Mean age was 63 ± 18 years. The proportion of men was clearly higher than that of women (77% vs 23%). Mean response time was 7 minutes 54 seconds, with a significant

difference ($p < 0.05$) between cases of death (8 mins. 12 secs) and 7-day survivors (5 mins. 53 secs). Reasons for activating BLS units are shown in Figure 2. Confirmed CPA represented only 6% of the cases and "unconscious patient" was the reason in 31%. In the remaining 63% of cases, the data given by the caller (alerter) did not allow presumption of CPA. In only 14.8% of 7-day survivors, the reason for the alert was suspected CPA (Figure 3).

Of all CPA patients attended by BLS units, 54.5% were defibrillated by BLS technicians. In total, reversion of the CPA haemodynamic situation was achieved in 106 cases (45.5%). In all cases, definitive transfer to hospital was performed by ALS units after first being attended by BLS units. Of these, 53.3% were alive at 6 hours after admission to hospital, 46.7% at 24 hours, and at 7 days, 27 (25% of all admissions and 11.6% of all the 233 CPA attended by BLS units).

Of the 27 7-day survivors of CPA in a public space, 17 showed perfect neurological state (grade 1 of the Glasgow-Pittsburg Scale) with complete recovery or were attended for other associated pathologies. This number of patients constituted 7.29% of the 233 CPA cases first attended by BLS units. Survival of these cases was significantly higher than that of CPA patients who only received cardiopulmonary resuscitation (CPR) without defibrillation (75% versus 25%, $p < 0.001$).

Discussion

The first conclusion relates to the initial hypothesis of the study: to demonstrate the effectiveness of implementing AED in all BLS units; previously demonstrated in diverse studies^{6,14,15,20-25,30} and confirmed in this, given the number of CPA




Figure 1. Annual distribution of the 233 cases of cardiopulmonary arrest attended by Basic Life Sport units in this 5-year study.




Figure 2. Data on reasons for activation. CV: cardiovascular. Pathol: pathology. CPA: cardiopulmonary arrest. Acc: accident. Intox: intoxication.

recoveries and survivors, as well as neurological state scores²⁵ since 17 of 27 (62.9%) 7-day survivors showed perfect neurological state. There is another even more relevant statistic supporting the implementation of AED in BLS units, which is the fact that the initial alert information did not predict the real pathology in the great majority of cases (94%), which was the main reason for sending out BLS units in the first place. This reveals the great utility of their being equipped with these devices at all times.

There is currently important lack of knowledge in the general population about recognising a CPA situation, which is essential as the first link in the Survival Chain²⁶. Thus the necessity for public information campaigns such as the Alert Program implemented by SAMUR-PC since 10 years ago, which trains an average of 10,000 citizens in recognition of emergency situations, the early alert system and initiation of the first measures for re-

suscitation³⁵. In the same line, recent studies such as that by Caffrey et al²⁷ highlight the efficacy and benefit in terms of survival of Public Access Defibrillation programs such as that developed in Madrid with collaboration between SAMUR-PC and the company Telefónica, with more than 100 AED installed in towers called “rescate cardiaco” (cardiac rescue) situated in busy areas of the city³⁵.

Of course, response time is a crucial factor when comparing results. De Maio et al¹² affirm that the objective of 8 minutes should be reduced to 7 minutes, which provides a further argument in favour of equipping BLS units with AED. In any emergency system, BLS units constitute the greatest percentage of ambulances, and therefore a higher number of adequately distributed AED, with greater potential proximity to CPA events and early defibrillation. Other recent studies confirm the benefit of initial use of defibrillators in terms of survival. Bunch et al²⁸ affirm that the survival of CPA patients after out-of-hospital FV increased by 12% following the implementation of an early defibrillation program. In studies by Hallstrom et al¹⁶ and Uriarte et al³¹ published in August 2004 in the New England Journal of Medicine, 30 of 130 CPA patients initially treated with AED survived, while in the group of patients only receiving initial CPR 15 of 107 CPA patients survived ($p = 0.03$; $RR = 2.0$).

Finally, the percentage of BLS-attended survivors without neurological deficit was 7.29%, just below that for non-presential CPA (cases occurring in the presence of health-care professionals were ruled out, which had the greatest percentage of survival) attended by SAMUR-PC and oscillating




Figure 3. Reasons for activation in the 27 CPA survivors at 7 days. Pathol: pathology. CPA: cardiopulmonary arrest. Intox: intoxication.

around 12%. Similar studies have reported admission survival rates of 2.72% (study by Uriarte et al³¹, published in EMERGENCIAS 2001) and global survival of 4.1%

in a study by Lim et al³² published in Resuscitation (without clarifying whether cases of presential CPA were also included). In the SAMUR-PC service, this figure is used as an indicator of management, and is considered to objectively reflect the general running of the service³³.

With adequate and continuous training, Emergency Technicians are capable of providing excellent instrumental BLS care of CPA patients CPA²⁰⁻²⁵, increasing the survival and neurological recovery rates of patients suffering CPA in an out-of-hospital setting. In our study we observed an increasing global number of cases with early defibrillation over the 5-year period, in accord with greater presence and competence of the BLS units in daily SAMUR-PC practice.

References

- 2005 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. Part 5 Electrical Therapies: Automated External Defibrillators, Defibrillation, Cardioversion, and Pacing. *Circulation* 2005;112(IV):35-46.
- 2005 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 8: Interdisciplinary topics. *Resuscitation* 2005;67:305-14.
- Perales Rodríguez de Viguri N, Jiménez Murillo L, González Díaz G, Alvarez Fernández JA, Medina Alvarez JC, Ortega Carnicer J. La desfibrilación temprana: conclusiones y recomendaciones del I Foro de Expertos en Desfibrilación Semiautomática. *Emergencias* 2002;14:328-35.
- Ayuso Baptista F, Jiménez Moral G, Fonseca del Pozo FJ, Ruiz Madruga M, Garijo Pérez A, Jiménez Corona J et al. Nuevos horizontes frente a la muerte súbita cardiaca: la desfibrilación externa semiautomática. *Emergencias* 2003;15:36-48.
- Lobatón CR, Varela-Portas Mariño J, Iglesias Vázquez JL, Rodríguez MDM. Results of the introduction of an automated external defibrillation programme for non-medical personnel in Galicia. *Resuscitation* 2003;58:329-3.
- Lobatón CR, Varela-Portas Mariño J, Iglesias Vázquez JL, Rodríguez MDM. Resultados de la implantación en Galicia de la desfibrilación semiautomática por personal no médico. *Emergencias* 2003;15:11-6.
- Acebedo Esteban FJ, Fernández González MT. Desfibrilación Semiautomática Realmente Precoc. [Carta]. *Emergencias* 1998;10:135-6.
- Wik L, Hansen TB, Fylling F, Steen T, Vaagenes P, Auestad BH, et al. Delaying defibrillation to give basic cardiopulmonary resuscitation to patients with out-of-hospital ventricular fibrillation: a randomized trial. *JAMA* 2003;289:1389-95.
- Smith KL, Cameron PA, Peeters A, Meyer AD, McNeil JJ. Automatic external defibrillators: changing the way we manage ventricular fibrillation. *Med J Aust* 2000;172:384-8.
- Cobb LA, Fahrenbruch CE, Walsh TR, Copass MK, Olsufka M, Breskin M, et al. Influence of cardiopulmonary resuscitation prior to defibrillation in patients with out-of-hospital ventricular fibrillation. *JAMA* 1999;281:1182-8.
- Lui JC. Evaluation of the use of automatic external defibrillation in out-of-hospital cardiac arrest in Hong Kong. *Resuscitation* 1999;41:113-9.
- De Maio VJ, Stiell IG, Wells GA, Spaite DW. Optimal defibrillation response intervals for maximum out-of-hospital cardiac arrest survival rates. *Ann Emerg Med* 2003;42:242-50. Comment in: *Ann Emerg Med* 2003 Aug;42:251-5.
- Nichol G, Stiell I, Laupacis A, Ba' Pham M, Valerie M, George W. A cumulative meta-analysis of the effectiveness of defibrillator capable emergency medical services for victims of out-of-hospital cardiac arrest. *Ann Emerg Med* 1999;34:517-25.
- Richardson LD, Kwun R, Mccburnie MA, Chason K. New approaches to out-of-hospital cardiac arrest. *Mt Sinai J Med* 2006;73:440-8.
- Cappato R, Curnis A, Marzollo P, Mascioli G, Bordonali T, Beretti S, et al. Prospective assessment of integrating the existing emergency medical system with automated external defibrillators fully operated by volunteers and laypersons for out-of-hospital cardiac arrest: the Brescia Early Defibrillation Study (BEDS). *Eur Heart J* 2006;27:508-9.
- Hallstrom AP, Ornato JP, Weisfeldt M. The Public Access Defibrillation Trial Investigators. Public-access defibrillation and survival after out-of-hospital cardiac arrest. *N Engl J Med* 2004;351:637-46.
- Stiell IG, Wells GA, DeMaio VJ, Spaite DW, Field BJ, Munkley DP, et al. Modifiable factors associated with improved cardiac arrest survival in a multicenter basic life support/defibrillation system: OPALS Study Phase I results. *Ann Emerg Med* 1999;33:44-50.
- Stiell IG, Wells GA, Field BJ, Spaite DW, De Maio VJ, Ward R, et al. Improved out-of-hospital cardiac arrest survival through the inexpensive optimization of an existing defibrillation program: OPALS study phase II. *JAMA* 1999;281:1175-81.
- Auble TE, Menegazzi JJ, Paris PM. Effect of out-of-hospital defibrillation by basic life support providers on cardiac arrest mortality: a meta-analysis. *Ann Emerg Med* 1995;25:642-8.
- Calle PA, Verbeke A, Vanhaute O, Van Acker P, Martens P, Buylaert W. The effect of semi-automatic external defibrillation by emergency medical technicians on survival after out-of-hospital cardiac arrest: an observational study in urban and rural areas in Belgium. *Acta Clin Belg* 1997;52:72-83.
- Ladwig KH, Schoefinius A, Danner R, Gurtler R, Herman R, Koeppel A, et al. Effects of early defibrillation by ambulance personnel on short- and long-term outcome of cardiac arrest survival: the Munich experiment. *Chest* 1997;112:1584-91.
- White RD, Asplin BR, Bugliosi TF, Hankins DG. High discharge survival rate after out-of-hospital ventricular fibrillation with rapid defibrillation by police and paramedics. *Ann Emerg Med* 1996;28:480-5.
- Mols P, Beaucarne E, Bruyninx J, Labruyere JP, De Myttenaere L, Naeije N, et al. Early defibrillation by EMTs: the Brussels experience. *Resuscitation* 1994;27:129-36.
- Sedgwick ML, Watson J, Dalziel K, Carrington DJ, Cobbe SM. Efficacy of out of hospital defibrillation by ambulance technicians using automated external defibrillators. The Heartstart Scotland Project. *Resuscitation* 1992;24:73-87.
- Acevedo Esteban FJ, Fernandez Gonzales MT, Suarez Bustamente RM, Rey Paterna P, Florez IC, Gilarranz Vaquero JL. Automated defibrillation performed by emergency medical technicians: the Madrid experience. *Resuscitation* 2000;43:155-7.
- Cummins RO, Ornato JP, Thies WH, Pepe PE. Improving survival from sudden cardiac arrest: the "chain of survival" concept. A statement for health professionals from the advanced Cardiac Life Support Subcommittee and the Emergency Cardiac Care Committee, American Heart Association. *Circulation* 1991;83:1832-47.
- Caffrey SL, Willoughy PJ, Pepe PE, Becker LB. Public Use of Automate external Defibrillators. *N Engl J Med* 2002;347:1242-7.
- Bunch TJ, White RD, Gersh BJ, Shen WK, Hammill SC, Packer DL. Outcomes and in-hospital treatment of out-of-hospital cardiac arrest patients resuscitated from ventricular fibrillation by early defibrillation. *Mayo Clin Proc* 2004;79:613-9.
- León Miranda MD, Gómez Jiménez FJ, Martín-Castro C, Cárdenas Cruz A, Olavarría Govantes L, Higuera Torres-Pucho J. Factores pronósticos de mortalidad en la parada cardiorrespiratoria extrahospitalaria. *Med Clin (Barc)* 2003;120:561-4.
- Pell JP, Sirel JM, Marsden AK, Ford I, Cobbe SM. Effect of reducing ambulance response times on deaths from out of hospital cardiac arrest: cohort study. *BMJ* 2001;322:1385-8.
- Uriarte E, Alonso D, Odriozola G, Royo I, Chocarro I, Alonso J. Supervivencia de la parada cardiorrespiratoria extrahospitalaria en Guipúzcoa: cuatro años de seguimiento. *Emergencias* 2001;13:381-6.
- Lim SH, Anantharaman V, Teo WS, Chan YH, Chee TS, Chua T. Results of the first five years of the prehospital automatic external defibrillation project in Singapore in the "Utstein style". *Resuscitation* 2005;64:49-57.
- Suárez Bustamante R, Corral Torres E, De Elías Hernández R, Moreno Martín JI, Bernardino Santos M, Gilarranz Vaquero JL. La Parada Cardiorrespiratoria como indicador de calidad de un Servicio de Emergencias Médicas. En: XV Congreso Nacional de la Sociedad Española de Medicina de Emergencias. Valencia. 2003.
- Sayre MR, Travers AH, Daya M, Greene HL, Salive ME, Vijayaraghavan K, et al. Measuring survival rates from sudden cardiac arrest: the elusive definition. *Resuscitation* 2004;62:25-34.
- Luna Parro S, Gacimartin Maroto E, Bernal del Valle D, Aguilar Fernández F, Palacio M, Corral Torres E. Columnas de Rescate Cardíaco. Desfibrilación Externa Automática al alcance de la Población General. En II Jornadas de Salud Pública de la Ciudad de Madrid.

Efectividad de la implementación de la desfibrilación semiautomática en la Unidades de Soporte Vital

Moreno Martín JL, Esquilas Sánchez O, Corral Torres E, Suárez Bustamanete RM, Vargas Román MI

Objetivo: Evaluar la importancia de la aplicación de la desfibrilación semiautomática en el tratamiento de paradas cardiorrespiratorias (PCR) no presenciadas, de cualquier etiología, atendidas inicialmente por unidades de Soporte Vital Básico (SVB), mediante el análisis de la supervivencia y valoración neurológica a los 7 días del suceso.

Método: Estudio prospectivo, descriptivo transversal/longitudinal, que analiza la supervivencia y el estado neurológico mediante la Escala de Glasgow-Pittsburg a los 7 días de la PCR de 233 pacientes atendidos durante un periodo de 5 años por el servicio urbano de emergencias extrahospitalarias de doble escalón de Madrid (SAMUR-PC).

Resultados: La supervivencia "ad integrum" tras una PCR no presenciada fue de 7,29%, muy próxima a los resultados globales de recuperación en nuestro servicio (12%), en los que se incluyen las PCR atendidas inicialmente por unidades de Soporte Vital Avanzado (SVA). Se constata una diferencia significativa entre los pacientes desfibrilados precozmente por SVB y aquéllos que por su ritmo inicial no lo habían sido. La demanda inicial de asistencia sólo hizo prever una PCR en el 37,7% de todos los casos.

Conclusiones: La primera respuesta a la PCR de origen no traumático con unidades de SVB con desfibrilador semiautomático (DEA) puede alcanzar niveles de supervivencia similares a la asistencia inicial por SVA. [Emergencias 2009;21:12-16]

Palabras clave: Soporte Vital Básico. Desfibrilación semiautomática. Técnicos de Emergencia. Parada cardiorrespiratoria. Supervivencia.